



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$= \frac{48}{\pi^6} \int_0^{\frac{1}{2}\pi} \int_0^\theta \int_0^\phi [(\pi-2\theta)(\pi-2\phi)(\pi-2\psi)] d\theta d\phi d\psi = \frac{1}{8}.$$

$$A = \frac{\frac{1}{2} \int_0^a \int_0^a \int_0^a \int_0^\pi \int_0^\pi [xysin \delta + yzs sin r - xzs sin(\delta+r)] x dx y dy z dz d\delta d r}{\int_0^a \int_0^a \int_0^a \int_0^\pi \int_0^\pi x dx y dy z dz d\delta d r}$$

$$= \frac{8}{\pi a^6} \int_0^a \int_0^a \int_0^a (x+z)xy^2z dx dy dz = \frac{8a^3}{9\pi}.$$

PROBLEMS FOR SOLUTION.

ALGEBRA.

357. Proposed by V. M. SPUNAR, M. and E. E., Chicago, Illinois.

Solve the system

$$\begin{aligned} \sqrt{x^2+a^2+b^2+c^2} &= \sqrt{y^2+b^2+c^2} + \sqrt{z^2+b^2+c^2}, \\ \sqrt{y^2+a^2+b^2+c^2} &= \sqrt{x^2+c^2+a^2} + \sqrt{z^2+c^2+a^2}, \\ \sqrt{z^2+a^2+b^2+c^2} &= \sqrt{x^2+a^2+b^2} + \sqrt{y^2+a^2+b^2}. \end{aligned}$$

358. Proposed by V. M. SPUNAR, M. and E. E., Chicago, Illinois.

Show that
$$\frac{n(n+1)\dots(n+m-1)}{m!} - n \frac{n(n+1)\dots(n+m-4)}{(m-3)!} + \frac{n(n-1)}{2!} \cdot \frac{n(n+1)\dots(n+m-7)}{(m-6)!} - \dots = 0, \text{ if } m > 2n; \text{ and } = 1 \text{ if } m = 2n.$$

359. Proposed by V. M. SPUNAR, M. and E. E., Chicago, Illinois.

Show when
$$1/(1-x)(1-x^3)(1-x^5)(1-x^7)\dots = (1+x)(1+x^2)(1+x^3)(1+x^4)\dots$$

GEOMETRY.

390. Proposed by PROF. R. C. ARCHIBALD, Brown University, Providence, R. I.

Find, geometrically and without introducing focal properties, the locus of the vertices of the conjugate parallelograms of an ellipse.

391. Proposed by W. J. GREENSTREET, M. A., Editor, The Mathematical Gazette, Stroud, England.

An ellipse is inscribed in the triangle of reference and has one focus at (sec A, sec A, sec C). Find the other focus and the sum of the squares of the axes of the ellipse.

392. Proposed by V. M. SPUNAR, M. and E. E., Chicago, Ill.

A tangent to a curve at any point P cuts the tangent and the normal at a fixed point O in the points M and N , and the rectangle $OMP'N$ is completed. Find the curve which is such that the triangle formed by the tangents at any three points P, Q, R is equal to the triangle formed by the corresponding points P', Q', R' .